

Insights from the Evidence Base

Dr Lynne Boddy & Dr Zoe Knowles



@LJMU_PAEx

Our agenda

- Physical activity: A complex behaviour
- Environments/opportunities to facilitate physical activity
- Measurement and evaluation of physical activity interventions
- Forest School
- Summary

Physical Activity: Definitions and guidance

Physical Activity:

“Any bodily movement produced by the skeletal muscles that substantially increases energy expenditure” Howley, 2001.


Sedentary behaviour [from the Latin *sedére* meaning ‘to sit’]:

“A distinct class of behaviours (e.g. sitting, lying) characterized by little physical movement and low energy expenditure” Tremblay *et al.*, 2011.

Chief Medical Officers guidelines:

<https://www.gov.uk/government/publications/uk-physical-activity-guidelines>

Physical Activity: A complex behaviour



Children's PA & Health

Insufficient physical activity increases disease risk in children

TABLE 3. Adjusted mean [SEM] clustered risk, CRF, and BMI Z-score by physical activity group controlling for gender, somatic maturation, and accelerometer wear time

Despite this, not enough children meet 60mins MVPA every day and levels of cardiorespiratory fitness have declined substantially in recent years.

TABLE 2. Adjusted mean [SEM] clustered risk, CRF, and BMI Z-score by physical activity group controlling for gender, somatic maturation, and accelerometer wear time

Variable	Inactive (n = 49)	Active (n = 18)	p-value
Clustered risk score	0.49 [0.53]	-1.74 [0.88] 97]	P = 0.04 P = 0.03 P = 0.14 P < 0.01

Variable	Unfit (n = 36)	Fit (n = 31)	P-value
Clustered Risk score	1.23 [0.60]	-1.66 [0.65]	P < 0.01
BMI Z-score	1.10 [0.15]	0.01 [0.16]	P < 0.01
Sedentary time (min)	530.77 [6.53]	492.33 [7.06]	P < 0.01
MVPA (min)	43.01 [4.03]	63.06 [4.36]	P < 0.01

cardiorespiratory fitness

Activity Measurement: How are children active?

- We are often concerned with *how active* children are.
- But, ***children's activity patterns*** influence how we measure PA within this population.
- Children's activity is sporadic, characterised by short bouts of high intensity activity, interspersed with periods of light activity and sedentary behaviour [Baquet et al., 2007]
- Children take part in a wide range of activities in a range of settings, so it is difficult to capture every activity bout.
 - Structured and un-structured
- Children have different preferences, barriers, facilitators and views about PA.
 - Difficult to capture using quantitative measures.

A combination of measures provides the best coverage.

When are children active?

- Boys are more active than girls at all ages and PA intensities [Sherar et al., 2007].
- Physical activity declines significantly with age [Trost et al., 2002].
- Children are usually more active on weekdays than weekends.
- Though most active children maintain activity levels at weekends [Fairclough et al., 2015].

Environments and Opportunities for children to be active

Schools & Physical Activity

- *Before school* – school is mandatory
- Infra-structure – teachers, facilities, curriculum
- *Recess periods* for 40+wk
Lots of opportunities for PA
- *PE & curriculum time* in the community, but
- >50% of daily MVPA accrued at school [Fairclough *et al.*, 2008;]
After school




Before/after-school: Active school travel


- Walking, cycling or combination with passive transport
- Active travel is simple, free and sustainable
- Decline in active travel over time
- Health Survey for England 2012:
 - Around 2/3s of children walk or cycle to /from school at least once a week
 - High school pupils spend ~17 min/day actively commuting.

How much PA does active school travel contribute?

- Cooper et al 2012: The PEACH Project




Children's recess PA & SB [Ridgers et al., 2012]




Physical activity promotion through PE

[Slingerland & Borghouts, 2011]



SB & inactivity in PE [Mersh & Fairclough, 2010]



After-school

- After-school period [up to 6.30pm] termed ‘critical hours’ for young people’s PA and SB [Atkin et al., 2008]
- Between $\frac{1}{3}$ and $\frac{1}{2}$ of young people’s PA may occur during this period [Tudor-Locke et al., 2006; Hager, 2006]
- Significant discrepancies in PA occur between the most and least active young people at this time [Fairclough et al., 2012]

How can we measure PA and its impact on children?

Subjective Assessments of Activity: Surveys

Strengths


- Low cost, can be used on a large scale.
- Give general information on physical activity levels.
- Also provide context (e.g. setting and type).
- Some useful surveys: PAQ-C & PAQ-A [Kowalski et al., 1997] Youth Activity Profile [Saint-Maurice et al., 2014].

Limitations

- Often provide scores rather than minutes of activity.
- Rely on participants understanding the question.
- Rely on accurate recall.
- Less precision than objective methods and usually overestimate activity.

Objective Assessment of Physical Activity: Accelerometers

- Small sensors that are commonly worn at the hip or wrist.
- Measure acceleration in 1-3 planes and convert these signals to minutes of activity.
- Reliable and valid for use in children.
- Provide detailed data.
 - Quantifies time spent across the activity intensity continuum.
 - Filter data by specific times, days, segments.
- Requires specialist software and training.
- Research decision making important in data processing.
- Monitors are expensive: ~£180 per device.




Qualitative research

- Formative
 - Compliance to device wear, e.g McCann et al [under review]; intervention design MackIntosh et al [2011]; Downs et al [2013]
- Evaluation of interventions
 - Ridgers et al, [2012] Knowles et al [2013]
- ‘Childs voice’
 - Focus groups, write and draw (WDST), creative methodologies

Qualitative research

Ridgers, N., Knowles, Z. & Sayers, J. [2012]



Qualitative research



Child as a ‘change agent’

- Child targeted intervention can create secondary reach
 - Child as a ‘change agent’
 - Ridgers [2012] Forest School
 - Houghton et al [2014] Active play intervention for Fathers/male carers and their pre school children
- Citizen science projects
 - PPI, public engagement agenda

Forest Schools

- Used the Physical Activity Questionnaire for Children [Kowalski et al., 1997] to examine PA levels pre and post intervention.
- Used accelerometers mid-intervention to examine the activity ‘dose’ from Forest Schools.
- Qualitative focus groups and write and draw techniques
- Results demonstrated:
 - Mean MVPA = 73.2 minutes on a Forest School day, 68.2 minutes on a PE school day, 67.6 minutes on a weekend day and 65.7 minutes on a normal school day.
 - Children participated in significantly more LPA on Forest School days in comparison to a regular school day and a weekend day.
 - Children participated in significantly more MPA Forest School day than on a regular non-PE school day.
 - However, children were also significantly more sedentary on Forest School days than weekends, but slightly less sedentary than on P.E. and normal school days.
 - No differences in pre-post intervention in PAQ score.

Austin et al [in review].

Forest School- Write and Draw

Flo

Sex: girl M/F
Age: 5

1. What I like about Forest School is...



Summary

- Evidence base regarding natural environments as PA environments is limited
- Interventions can be designed to promote PA amongst other benefits and create reach beyond the target beneficiary
- Measurement needs intervention and population appropriate methods
- Commitment to robust evaluation within the planning stages
 - Research partners

References

- Atkin, A. J., Gorely, T., Biddle, S. J. H., Marshall, S. J., & Cameron, N. (2008). Critical hours: Physical activity and sedentary behavior of adolescents after school. *Pediatric Exercise Science*, 20, 446-456.
- Austin, C., Boddy, L.M., Sayers, J & Knowles, Z. (Under review). Exploring the impact of Forest School: Physical activity and child led evaluation of the programme. *Childrens Geographies*
- Baquet, G., Stratton, G., Van Praagh, E., & Berthoin, S. (2007). Improving physical activity assessment in prepubertal children with high-frequency accelerometry monitoring: A methodological issue. *Preventive Medicine*, 44, 143-147.
- Cooper, A. R., Jago, R., Southward, E. F., & Page, A. S. (2012). Active Travel and Physical Activity across the School Transition: The PEACH Project. *Med Sci Sports Exerc*, 44(10), 1890-1897.
- Dollman, J., Okely, A. D., Hardy, L., Timperio, A., Salmon, J., & Hills, A. P. (2009). A hitchhiker's guide to assessing young people's physical activity: Deciding what method to use. *J Sci Med Sport*, 12(5), 518-525.
- Downs S., Knowles Z., Fairclough S.J., Stratton G. & Boddy, L.M. (2013). Exploring opportunities available and perceived barriers to physical activity engagement in children and young people with Down syndrome. *European Journal of Special Needs Education*. 28, 270-287
- Fairclough, S. J., Beighle, A., Erwin, H., & Ridgers, N. D. (2012). School day segmented physical activity patterns of high and low active children. *BMC Public Health*, 12(1), 406.
- Fairclough, S.J., Boddy, L.M., Mackintosh, K.A., Valencia-Peris, A., Ramirez-Rico, E. (2015) Weekday and weekend sedentary time and physical activity in differentially active children. *Journal of Science and Medicine in Sport*. 18(4): 444-449.
- Hager, R. L. (2006). Television viewing and physical activity in children. *Journal of Adolescent Health*, 39, 656-661.
- Houghton, L., O Dwyer, M. V., Foweather, L., Watson, P., Alford, S., & Knowles, Z. R. (2014). An impact and feasibility evaluation of a six-week (nine hour) active play intervention on fathers' engagement with their preschool children. *Early Child Development and Care*. 185 (2) 244-266.
- Kowalski KC, Crocker PRE, Faulkner RA. (1997) Validation of the physical activity questionnaire for older children. *Pediatric Exercise Science*, 9: 174-186.
- Knowles, Z., Parnell, D., Ridgers, N. & Stratton, G. (2013) Learning from the experts: Exploring playground experience and activities using a write and draw technique. *Journal of Physical Activity & Health*, 10, 406-415

References

- McCann, D., Knowles, Z., Fairclough, S & Graves, L. (Under review). Strategies to encourage accelerometer wear in children and young people. Qualitative Review in Sport and Exercise
- Mersh, R., & Fairclough, S. J. (2010). Physical activity, lesson context and teacher behaviours within the revised English National Curriculum for Physical Education: A case study of one school. European Physical Education Review, 16(1), 29-45.
- Mackintosh, K., Knowles, Z., Ridgers, N., & Fairclough, S. (2011). Using formative research to develop physical activity. CHANGE!: A curriculum-based health education intervention. BMC Public Health, 11, 831
- Ridgers, N., Knowles, Z. & Sayers, J. (2012). Play in the Natural Environment: A Child-Focused Evaluation of Forest School. Children's Geographies, 10, 55-71
- Ridgers, N. D., Timperio, A., Crawford, D., & Salmon, J. (2012). Five-year changes in school recess and lunchtime and the contribution to children's daily physical activity. British Journal of Sports Medicine, 46(10), 741-746
- Saint-Maurice PF, Welk GJ, Beyler NK, Bartee RT, Heelan KA. (2014) Calibration of self-report tools for physical activity research: the Physical Activity Questionnaire. BMC Public Health, 14P 461.
- Sherar, L. B., Esliger, D. W., Baxter-Jones, A. D. G., & Tremblay, M. S. (2007). Age and gender differences in youth physical activity: Does physical maturity matter? Medicine and Science in Sports and Exercise, 39, 830-835.
- Slingerland, M., & Borghouts, L. (2011). Direct and indirect influence of physical education-based interventions on physical activity: a review. [Review]. Journal of physical activity & health, 8(6), 866-878.
- The NHS InformationCentre. (2009). Health Survey for England 2008. Physical activity and fitness (Vol. 1). London: The Health and Social Care Information Centre.
- Trost, S. G., Rosenkranz, R. R., & Dzewaltowski, D. A. (2008). Physical activity levels among children attending after-school programs. Medicine and Science in Sports and Exercise, 40(4), 622-629.
- Trost, S. G., Pate, R. R., Sallis, J. F., Freedson, P. S., Taylor, W. C., Dowda, M., & Sirard, J. (2002). Age and gender differences in objectively measured physical activity in youth. Medicine and Science in Sports and Exercise, 34(2), 350-355.
- Tudor-Locke, C., Lee, S. M., Morgan, C. F., Beighle, A., & Pangrazi, R. P. (2006). Children's pedometer-determined physical activity during the segmented school day. Medicine and Science in Sports and Exercise, 38, 1732-1738.